

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 754 178 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

15.01.2003 Bulletin 2003/03

(51) Int Cl.⁷: **C07D 219/04**, G01N 33/533,
G01N 33/58, C07J 43/00

(21) Application number: **95913298.6**

(86) International application number:
PCT/IB95/00244

(22) Date of filing: **06.04.1995**

(87) International publication number:
WO 95/027702 (19.10.1995 Gazette 1995/45)

(54) **NOVEL FUNCTIONALIZED HYDROPHILIC ACRIDINIUM ESTERS**

FUNKTIONALISIERTE HYDROPHILE AKRIDINIUM ESTER

NOUVEAUX ESTERS D'ACRIDINIUM HYDROPHILES FONCTIONNALISES

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB IT LI

(30) Priority: **08.04.1994 US 225165**

(43) Date of publication of application:
22.01.1997 Bulletin 1997/04

(60) Divisional application:
99203889.3 / 0 982 298

(73) Proprietor: **Bayer Corporation**
East Walpole, MA 02032 (US)

(72) Inventors:
• **LAW, Say-Jong**
Westwood, MA 02090 (US)
• **SOTIRIOU-LEVENTIS, Chariklia**
Rolla, MO 65401 (US)
• **NATRAJAN, Anand**
Manchester, NH 03104 (US)
• **JIANG, Qingping**
Northborough, MA 01532 (US)

- **CONNOLLY, Peter, B.**
Walpole, MA 02081 (US)
- **KILROY, John, P.**
Boston, MA 02116 (US)
- **MCCUDDEN, Constance, R.**
Brookline, MA 02146 (US)
- **TIRRELL, Stephen, M.**
Franklin, MA 02038 (US)

(74) Representative: **Froud, Clive et al**
Elkington and Fife
Prospect House
8 Pembroke Road
Sevenoaks, Kent TN13 1XR (GB)

(56) References cited:
EP-A- 0 082 636 **EP-A- 0 263 657**
EP-A- 0 273 115 **EP-A- 0 353 971**
EP-A- 0 361 817 **EP-A- 0 609 885**

Remarks:

The file contains technical information submitted
after the application was filed and not included in this
specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 754 178 B1

DescriptionField of the Invention

[0001] The present invention relates to a novel method for detection of an analyte. The present invention relates to the detection of an analyte using acridinium esters as chemiluminescent markers which can be encapsulated within liposome vesicles without significant leakage of the esters from the vesicles. The present invention also relates to the synthesis and use of novel functionalized hydrophilic acridinium esters which are useful as chemiluminescent labels, and surprisingly give a much higher quantum yield than prior acridinium ester compounds. The present invention also relates to novel hydrophilic acridinium esters which may be used to form direct and indirect conjugates. The present invention also relates to novel conjugates formed from such functionalized hydrophilic acridinium ester compounds. The present invention further relates to assays utilizing these novel functionalized hydrophilic acridinium esters and conjugates thereof. The instant invention relates to immunoassay using the compounds of the instant invention.

Background of the Invention

[0002] The use of acridinium esters as chemiluminescent labels in clinical assays is known. For example, EP-A-00 82 636 describes the use of an aryl acridinium ester activated with an N-hydroxy-succinimidyl moiety as a chemiluminescent label in immunoassays. US-A-4,745,181 and US-A-4,918,192 (EP-A-0263657); US-A-5,241,070; and US-A-5,663,074 describe polysubstituted aryl acridinium esters (PAAE) which are useful in immunoassays and nucleic acid hybridization assays. US-A-5,227,489 and US-A-5,449,55 (EP-A-0353971) describe hydrophilic polysubstituted aryl acridinium esters and lumisome conjugates thereof useful in clinical assays, particularly those assays involving liposomes.

[0003] Previous methods for the synthesis of 2',6'-Dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl-10-Methyl-9-acridinecarboxylate Methylsulfate (DMAE-NHS) as described in US -A-4,745,181 and EP-A-0263657, require the use of a phenoxy group substituted with a benzyloxycarbonyl group as an intermediate to form the acridine ester via a long synthetic pathway. It is desirable to develop new and efficient methods of synthesizing the useful acridinium ester labels of the instant invention. The unexpected ability to form 2'6'-dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 9-acridinecarboxylate (DMAeE-NHS) by the simplified procedure of combining a solution of succinimidyl 3,5-dimethyl-4-hydroxybenzoate and 4-dimethylaminopyridine with 9-acridinecarbonyl chloride hydrochloride, was not readily predictable in view of the coexistence of two reactive leaving groups in the same reaction, an acid chloride from 9-acridinecarbonyl chloride and a succinimidyl ester from succinimidyl 3,5-dimethyl-4-hydroxybenzoate. For example, in the case where N-succinimidyl 3(4-hydroxyphenyl)-propionate is reacted with the 9-acridinecarbonyl chloride referred to above, the condensation can be carried out under mild conditions (approximately room temperature) due to the absence of the two methyl groups in the ortho positions. (See for example US-A-4,946,958, cols. 4 - 5, where the reactant does not contain the ortho-substituted methyl groups.) On the other hand, when the two methyl groups are present, much more drastic conditions (100 degrees C. for 2 hours) are required for the condensation reaction to take place, due to the steric hindrance caused by the methyl groups. It should be noted, though, that there is a benefit provided by the presence of the methyl groups, namely the added stability of the resulting acridinium ester, as discussed, for example, in US-A-4,745,181.

[0004] The lipophilic nature of the prior art acridinium esters and other chemiluminescent compounds render them unsuitable for encapsulation within liposomes because of their rapid leakage through the liposome wall. Additionally, the limited water solubility of prior art acridinium esters and other chemiluminescent compounds only allow the encapsulation of a few marker molecules per liposome vesicle, resulting in relatively low signal amplification.

[0005] The novel functionalized hydrophilic acridinium esters of the instant invention are useful in immunoassay, yield results that are superior in sensitivity to prior methods, and do not need to use dangerous radiolabels, or organic enzyme/substrates. The use of chemiluminescent labels of this type resulted in an unexpected improvement over prior methods and can lead to the functional improvement of previously inoperable or inaccurate assay methodologies. The novel discovery that hydrophilic acridinium esters of the instant invention could be used to label biomolecules and compounds directly without reducing the solubility of the complex allows for many applications in the field of immunoassay. The novel compounds of the instant invention will allow for sensitive immunoassay without the need of excessive use of reagents in the reaction mixture, and are thus designed to reduce non-specific interactions, while not interfering with desired specific interactions.

[0006] Another unexpected benefit of having an PAAE carrying a hydrophilic moiety, particularly as a substituent at the nitrogen atom of the acridinium nucleus, is the significant improvement of the chemiluminescent quantum yield relative to that of acridinium esters which simply have an alkyl group (US-A-4,745,181) or carboxymethyl group (G. Zomer et al., 1989, *Anal. Chem. Acta* **227**: 11-19) substituted at the same position.

[0007] Reference may also be made to the following: EP-A-0361817, which discloses nucleophilic polysubstituted

aryl acridinium esters, conjugates, production and uses thereof; EP-A-0273115, which discloses chemiluminescent acridinium and phenanthridinium salts; and EP-A-0609885, which discloses acridinium compounds and conjugates thereof.

[0008] The area of immunoassay is well developed, and it is the discovery of unique labeling systems that can spur a quantum advancement in the field of immunoassay. In the field of immunoassay it is desirable to have assays which are highly specific, and highly sensitive to low concentrations, and yet still yield detectable and distinguishable signals for measurement. The instant invention allows for the specific detection of analytes at low concentration, without loss of solubility. The instant invention provides a means for making many specific labeling agents which can be used to detect low levels of analytes. The instant invention also teaches one with ordinary skill in the art, a novel and useful means of performing immunoassay that lends itself readily to automation and commercialization as assay kits and reagent kits.

[0009] The application of the novel acridinium esters of the instant invention in conjugates with bioactive proteins such as avidin, antibodies, DNA binding proteins, histones, and ribosomes, and others, is made possible by the hydrophilic properties of the synthesized compounds. These novel acridinium esters are also useful for the labeling of isolated, or intact, RNA, DNA, proteins, peptides, inactivated protein, neurotransmitters, hormones, viruses, viral antigens, bacteria, bacterial antigens, toxins, cytokines, antibody fragments, receptor proteins and other such targets both *in vitro*, and *in vivo*. The sampling of tissue samples, serum samples or other such biological samples, and the detection of previously difficult to measure specific analytes in rapid assays, are made possible by the compositions and methods of the instant invention.

[0010] one specific application of the instant invention is in the field of pathogen detection. While there have been many assays developed to detect viral pathogens, such as Rubella, the instant invention provides a unique and highly sensitive assay which is unexpectedly superior to previous chemiluminescent methods. The instant methods are comparable with conventional assays in sensitivity yet are more efficient in time required to perform the assay.


[0011] Another application of the instant invention is in the area of detection of hormones or haptens or other small biologically active molecules in biological samples. Where the hormone or hapten levels are low, and transient, it is useful to have a sensitive and rapid method of testing for, and measuring the levels of, hormones or haptens in such biological samples. One of the possible examples is the detection of steroid hormone levels such as testosterone. Detection of testosterone using homologous and heterologous hapten conjugates is known. In this methodology, antibodies are generated to one form of testosterone immunogen. Then said antibodies are used to detect testosterone in a sample, while a conjugated competitive hapten tracer, that is of a different form than the immunogen, is added. Homologous assays are burdened with unacceptably high cross-reactivity such as when using C4-testosterone-B-gal conjugate haptens, and T-3-O-CMO-glucoamylase conjugates. Somewhat lower crossreactivity has been obtained with 11a-substituted hapten-HRP, and T-3-O-CMO hapten-penicillinase (Rao et al., 1992, Steroids 57: 154-162). The instant invention provides for superior detection over prior heterologous hapten assay in that the hydrophilic acridinium ester labels of the instant invention contribute to reduced non-specific interactions.

[0012] Accordingly, it is the purpose of the instant invention to provide novel functionalized hydrophilic acridinium esters and conjugates thereof for use as chemiluminescent tracers. It is also a purpose of the instant invention to provide for novel methods of detecting an analyte using functionalized hydrophilic acridinium esters and conjugates thereof. It is also a purpose of the instant invention to provide a new improved synthetic procedure for the efficient production of acridine esters.

Summary of the Invention

[0013] Novel acridinium esters that are useful, either alone or when incorporated into liposomes, as chemiluminescent agents in binding assays (e.g., immunoassays and gene probe assays) with improved sensitivity are disclosed. In addition, the synthesis of these esters and their use in assays for detecting an analyte is described. In particular, assays for testosterone and the Rubella virus are disclosed.

[0014] In one embodiment, the present invention provides a hydrophilic acridinium ester characterised in that it corresponds to the following general formula:



wherein R_1 represents sulfopropyl or sulfoethyl; R_2 represents hydrogen, methoxy, ethoxy, nitro or halogen; R_5 and R_7 represent hydrogen; R_4 and R_3 independently represent methyl, ethyl or isopropyl; and $-R_6$ represents N-succinimidylloxycarbonyl;

and

wherein R_5 , R_6 and R_7 substituent positions on the phenoxy ring are interchangeable.

[0015] In another embodiment, the present invention provides such an acridinium ester covalently bonded directly or via a bifunctional linker selected from hexyl-1,6-diamine, ethylene diamine or aminocaproic acid to a small biologically active molecule or a macro-biomolecule.

[0016] In a further embodiment, the present invention provides a method for measuring the amount of analyte in a sample characterised in that it comprises:

(a) detecting an analyte specifically and in proportion to its concentration in a sample;

and

(b) measuring a signal which is generated by the acridinium ester that is directly or indirectly proportional to the concentration of an analyte in a sample.

[0017] Having indicated the scope of the present invention, it will now be further described and illustrated in more general terms.

Brief Description of the Drawings

[0018] FIG. 1 is a diagram illustrating the synthesis of DMAE-NHS (2',6'-dimethyl-4'-(N-succinimidylloxycarbonyl) phenyl 10-methyl-9-acridiniumcarboxylate methosulfate; not claimed)

[0019] FIG.2 is a diagram illustrating the old method of synthesis of DMAE-NHS.

[0020] FIG. 3 is a standard curve for a testosterone chemiluminescence immunoassay using NSP-DMAE-HD-19-HC-MT (N-sulfopropyl dimethyl acridinium ester conjugate with 19-hydroxy-19-carboxymethyl-testosterone cross-linked with *n*-hexyl-1,6-diamine) of the instant invention as the tracer.

[0021] FIG. 4 is a comparison of binding curves for heterologous and homologous tracers in a chemiluminescence immunoassay for testosterone using the compounds of the instant invention.

[0022] The figure also shows a comparison between using hydrophilic NSP-DMAE vs. non-hydrophilic DMAE acridinium esters.

(ED₅₀ - analyte dose that corresponds to 50% displacement of the maximum tracer binding, or bound signal at particular dose/bound signal at 0 dose = 50%;

- - NSP-DMAE-HD-19-HCMT, a NSP-DMAE/19-HCMT conjugate cross-linked with *n*-hexyl-1,6-diamine;
- ▽ - NSP-DMAE-HD-19-CVNT, a NSP-DMAE conjugate with 19-carboxyvinyl-nor-testosterone cross-linked with *n*-hexyl-1,6-diamine;
- - DMAE-HD-19-HCMT, a DMAE/19-HCMT conjugate cross-linked with *n*-hexyl-1,6-diamine;
- ▼ - NSP-DMAE-HD-19-CMET, a NSP-DMAE conjugate with testosterone-19-CME cross-linked with *n*-hexyl-1,6-diamine.)

[0023] FIG. 5 is a plot showing the correlation of results from an ACS testosterone immunoassay using the compounds of the instant invention with DPC Coat-a-Count radioimmunoassay (Diagnostic Products Corp., Los Angeles, CA).

[0024] FIG. 6 is a bar graph showing a population study comparing the positive to negative ratio of assay results using the compounds of the instant invention as compared with DMAE rubella assay.

(■ - conjugate of DMAE with Rubella;
□ - conjugate of N-sulfopropyl DMAE with Rubella.)

[0025] FIG. 7 shows the assay architecture for an ACS Testosterone assay, which is run on the Ciba Corning ACS-180 instrument.

[0026] FIG. 8 shows the assay architecture for an ACS Estradiol assay, which is run on the Ciba Corning ACS-180 instrument.

[0027] FIG. 9 shows the assay architecture for an ACS Specific IgE assay, which is run on the Ciba Corning ACS-180 instrument.

Description of the Invention

[0028] The following terms as used in the specification and claims shall have the following meanings:

Analyte - the compound or composition to be measured which can be a ligand that is mono- or polyeptopic, antigenic, or haptenic. The analyte can be a piece of DNA or RNA. The analyte can be found in liquid or on solid support.

Conjugates - the combination of chemiluminescent compound with a second compound or molecule. Such conjugates can have many chemiluminescent molecules per second compound.

Indirect Conjugate - the combination of chemiluminescent compound with a second compound or molecule. The combined chemiluminescent compound with a second compound or molecule is then capable of further combining with a third compound or molecule. Such conjugates can have many chemiluminescent molecules per complex.

Homologous - the same compound linked by the same linker, at the same position.

Heterologous - the compound linked by a different linker at a different position, a different linker at the same position, the same linker at a different position.

Linker - a chemical bridge, tie, connection between two compounds or molecules; bifunctional linker. The binding of a linker to one compound can allow for a reactive site to be made available for further reaction.

Hapten - an incomplete antigen, incapable by itself to provoke an immune response, but when suitably attached to another molecule becomes capable of producing antibodies which will specifically recognize the hapten.

[0029] The hydrophilic PAAE of the instant invention, with an additional electrophilic or leaving group is useful for conjugation to many targets. The hydrophilic moiety of such a PAAE (comprising an alkyl spacer carrying a highly ionizable group such as sulfonate) is responsible for enhancing the aqueous solubility, hence reducing the hydrophobic nature of the tracers, particularly those formed from conjugating the PAAE with small organic biomolecules such as steroids, therapeutic drugs, and other controlled chemical substances. When applied in labelling macro-biomolecules such as proteins, nucleic acids, and polysaccharides, the hydrophilic PAAE permits the increase in incorporation ratio (defined as number of PAAE molecules covalently attached to a single macro-biomolecule) without any resulting solubility problems. Thus the increase in specific activity and washability of the tracer would be a factor in enhancing the binding assay sensitivity and improved signal/noise ratio. The lipophilic nature of the prior art acridinium esters and other chemiluminescent compounds render them difficult to use when conjugated to certain biological compounds, since there was a significant amount of non-specific binding due to the hydrophobic nature of the tracer conjugate. Thus the prior art acridinium esters and other chemiluminescent compounds were not efficient for use when detection required fine distinction between signal from specific vs. non-specific binding, and resulted in an increased risk of false positive or false.

[0030] It was unexpectedly found that PAAE with a substituted hydrophilic moiety (in particular 2',6'-Dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 10-(3'-Sulfopropyl)-acridinium-9-carboxylate (NSP-DMAE-NHS)) is capable of generating about 1.5 times higher chemiluminescent quantum yield as compared to the corresponding PAAE with simple alkyl substitution at the same position (i.e. 2',6'-Dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 10-Methyl-9-acridine-carboxylate Methylsulfate (DMAE-NHS)). (See Table I). The increase in chemiluminescent quantum yield of PAAE is a very desirable and essential factor in improving the signal to noise (S/N) ratio of the chemiluminescent tracers and the sensitivity of chemiluminescent binding assays and detection schemes. In particular, where the signal enhancement of the tracer is mediated through multiple PAAE labelling of a conjugate of ligand/carrier biomolecule such as a protein,

a 1.5 fold or greater improvement in PAAE chemiluminescent quantum yield will mean that less PARE is required to be bound to achieve the same signal amplitude. Also, if the same levels of binding of PARE to carrier conjugates is achieved, the detection of lower concentrations of an analyte may be possible.

TABLE I

Relative Quantum Yield of DMAE-NHS ⁽¹⁾ vs. NSP-DMAE-NHS ⁽²⁾				
AE ⁽³⁾ Type	Mol. Wt.	RLU/4.2 fmol AE		R L U ⁽⁴⁾ Ratio
DMAE-NHS	595		690,187 644,587 <u>743,587</u> mean 692,797 SD ⁽⁵⁾ 49,551 % C.V. ⁽⁶⁾ 7.2	1.0
NSP-DMAE-NHS	591		1,122,017 1,204,017 <u>1,122,017</u> mean 1,149,350 SD 47,343 % C.V. 4.1	1.7

(1) 2',6'-dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 10-methyl-9-acridiniumcarboxylate methosulfate

(2) 2',6'-dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 10-(3'-sulfopropyl)-acridinium-9-carboxylate

(3) acridinium ester

(4) relative light units

(5) standard deviation

(6) percent coefficient of variation

[0031] In table I, DMAE-NHS was dissolved in acetonitrile, while NSP-DMAE-NHS in 50% (v/v) acetonitrile in water to generate stock solutions of 1.0 mg/ml. Serial dilutions were carried out to 10^{-7} with flashing buffer (10mM Phosphate, 150 mM NaCl, 0.1% BSA, 0.05% NaN₃, pH 8.0) diluent. The dilution method consists of three 100-fold dilution steps using 50 uL transfer volumes and on 10-fold dilution step using 100 uL transfer volume. The light emission of the samples were determined on Ciba Corning Diagnostics MLA1 Analyzer under standard conditions (2 sec light collection time) using 25 uL of the 10^{-7} dilution solutions.

[0032] There are many modifications which one with ordinary skill in the art will recognize are within the scope of the instant specification. A number of spacers can be attached which will facilitate the connection of the hydrophilic group to the nitrogen atom at the acridinium nucleus, and an electrophile or leaving group that will facilitate the covalent attachment of a hydrophilic PAAE to other compounds or biomolecules.


[0033] The novel acridinium esters of the present invention are highly soluble in water and can be encapsulated in liposomes at high concentrations. Once inside liposomes at high concentrations, the novel acridinium esters remain encapsulated for extended periods of time and do not leak appreciably.

[0034] It will be appreciated that while the novel acridinium esters of the present invention are useful for encapsulation within liposomes, the novel acridinium esters of the present invention are also useful in other applications where acridinium esters are utilized, such as labeling ligands or analytes (such as antigens); labeling the specific binding partners of ligands or analytes (such as the corresponding antibodies); or labeling nucleic acids and molecules comprising nucleic acids.


[0035] In particular, the novel functionalized hydrophilic acridinium esters of the present invention will be useful for labeling biological materials for chemiluminescent assays where it is preferable to reduce the non-specific binding of tracer to biological material. The novel functionalized hydrophilic acridinium esters of the present invention will allow for a reduction in signal/noise (S/N ratio) interference, which will allow for greater accuracy in making measurements and for diagnostic assays, since the hydrophilic nature of the novel compounds of the instant invention will be less prone to non-specific interactions in comparison with other chemiluminescent compounds.

[0036] Synthesis of acridinium esters have been described as in US-A-4,745,181 and the process is shown in FIG. 2. This method involved the use of a phenoxy group substituted with a benzyloxycarbonyl group to form the acridine ester. The desired succinimidylloxycarbonyl was then introduced through the removal of the benzyl protecting group and the condensation of the product with N-hydroxysuccinimide in the presence of dicyclohexylcarbodiimide.

[0037] In addition, the instant invention provides for a new and improved method of synthesis. This method takes advantage of the unexpected discovery that the acridine nucleus, under properly selected conditions, can be directly condensed with the phenoxy group carrying succinimidyloxycarbonyl, without any problem, this is illustrated in FIG. 1. By using this new and simpler process, two synthetic steps of the older process are eliminated, for the preparation of hydrophilic PAAE with the proper electrophilic or leaving group. The generalized reaction scheme of Figure 1 also applies to the preparation of substituted acridine acid chlorides, such as



and N-hydroxysuccinimide ester of substituted p-hydroxybenzoates, such as




where R2, R5, R7 are hydrogen, halide, nitro, -R, -OR, -CN, -NHCOR, -COR, -COOR, or -CONHR; where R is alkyl, alkenyl, alkynyl, or aralkyl, and where R4 & R3 are R, and R is as defined above.


[0038] Thus the instant invention provides for hydrophilic PAAE with an additional electrophilic or leaving group, which are useful for conjugating with mono- or poly-nucleophilic compounds or biomolecules. The instant invention thus provides for the covalent attachment of the functionalized hydrophilic PAAE with nucleophilic compounds such as hexyl-1,6-diamine, ethylene diamine, and aminocaproic acid. The instant invention also provides for the covalent attachment of the functionalized hydrophilic PAAE with biomolecules such as aminoglycosides, proteins, viral antigens, peptides, and aminoalkyl-functionalized oligonucleotides. These conjugates will be useful for sensitive immunoassays and nucleic acid binding assays.


[0039] An example of a useful sensitive chemiluminescent immunoassay utilizing the compounds and conjugates of the instant invention is one which can detect specific steroid hormones in serum samples. An improved assay using the conjugates and methods of the instant invention is a speedy assay for testosterone. This assay combines the use of a highly specific antibody raised to testosterone and a heterologous tracer formed by conjugating a novel C19-mono- or C19-bi-substituted testosterone with a functionalized, hydrophilic PAAE. The benefit of using the heterologous chemiluminescent tracer is observed through the improvement of testosterone assay sensitivity to 0.1 ng/mL, which cannot be otherwise achieved rapidly (ie. about 5 min.) with the use of the corresponding homologous tracer, or using hydrophobic PAAE.

[0040] Thus the instant invention also provides for the utilization of functionalized hydrophilic PAAE which were formed as conjugates by novel C19-C-linked type of testosterone derivitization (as described in the Examples to follow). These novel heterologous tracers can be formed, optionally with further olefinic linkage at C19, or additional heteroatom-linked substitution at C19, such as the following C19 - O linked derivative of testosterone (NSP-DMAE-HD-19-CMET).



[0041] Thus the instant invention also provides for testosterone derivatives of the following structures. (Seven examples are shown below, along with a template identifying the numbering system used for nomenclature.)






wherein A is a functionalized group selected from the group consisting of -OH, -N(R₁)₂, -NHR₁, nitro, halide, -OR₁, -OCOR₁, -Tosylate, -mesylate, -azide, -isocyanate, -thioisocyanate, -SR₁, nitro, -SH, -SSR₁, -phosphonate, -sulfonate, -NHCOOR₁, -NHCOR₁, -NHCONH₂, hydrazines, cyanide, and -R₁; where R₁ is an alkyl, alkenyl, alkynyl, aralkyl containing up to 10 carbons and up to 5 heteroatoms; where B is -(CH₂)_n where n = 1-4 or -C≡; when B is -C≡, A is preferably omitted; where X is a carboxylate, -COOR₂, -CONHR₂, (where R₂ is an alkyl, alkenyl, alkynyl, aryl, aralkyl containing up to 15 carbons and up to 5 heteroatoms), or a carbonyl attached with proper leaving groups, including but not limited to halide, N-hydroxysuccinimide, and imidazole.

[0042] The enhanced utility of the hydrophilic PAEE of the instant invention in immunoassay of therapeutic drugs or hormones such as testosterone illustrates one aspect of the invention embodied in the instant specification. Specifically, the use of testosterone derivatives wherein the C19-B linked bridges (as shown in the above figure) are C19-C and an important element to the effective use of the compounds of the instant invention. The use of a C19-C linkage has been found to balance the reduction of tracer binding to the specific anti-testosterone antibody, and the ease of its displacement from antibody by competing sample testosterone.


[0043] A preferred embodiment of the testosterone tracers of the instant invention to be utilized for conjugation to the functionalized hydrophilic PAEE, via bifunctional crosslinkers, encompass the forms represented by formula I above. More specifically, it is preferred that A is hydroxy, B is -CH₂-, and X is carboxylate.

[0044] These compounds are novel, being distinguishable from those known from the prior art. For example, those compounds cited in Rao (US-A-4,197,286) contain C19 - o (i.e., C19 to oxygen) linkages.

[0045] Using the teachings of the instant specification one with ordinary skill in the art will recognize the use of bifunctional linkers for conjugation. A preferred embodiment encompasses a functionalized hydrophilic PAEE is attached to n-hexyl-1,6-diamine at one end through the formation of amide linkage, and subsequently to the target biomolecule, such as the testosterone derivative by the formation of another amide linkage with the other end, as exemplified by the following (NSP-DMAE-HD-19-HCMT).



[0046] Another preferred embodiment of the testosterone tracer is a form of the compound of formula I above, where A is omitted, B is -CH= forming an olefinic linkage with C19, and X is carboxylate, as shown below (NSP-DMAE-HD-19-CVNT).



[0047] A further aspect of the instant invention encompasses the unexpected improved results of using functionalized hydrophilic PAAE for labeling viral antigen as utilized in various infectious disease chemiluminescent immunoassays. For example, in an IgG capture assay format, where mouse anti-human IgGFc immobilized solid phase and viral antigen tracer are the reagents used for assaying anti-viral antibodies in the serum sample, it was found that great improvement in the signal to noise ratio of the assay can be achieved when a functionalized hydrophilic PAAE is used in place of the conventional DMAE-NHS in preparing the viral antigen tracer. These results most likely occur via better binding activity resulting in enhanced immunocomplex formation between tracer and target antibodies, and lowered non-specific binding due to the use of hydrophilic acridinium ester tags.

[0048] Thus one with ordinary skill in the art could take the teachings of the instant invention and apply them to the detection of a variety of infectious agents or pathogens. These infectious agents such as Rubella, Hepatitis of various classes and subtypes, Toxoplasma, Cytomegalovirus (CMV), HIV, chlamydia to name only a few, are targets for early detection which will require assays of great sensitivity and low incidence of false positive indication.

[0049] A typical architecture for such an assay consists of, but is not limited to, three components, a tracer, a sample containing the analyte, and a means for partitioning bound from unbound analyte. For example, such assays could combine the functionalized hydrophilic PAAE with suitable antigen of or specific binder for an analyte to form a tracer, incubation with a biological sample (containing the analyte), and capture of the tracer by a solid phase immobilized antibody. Other architectures will become practical with the application of the teachings of the instant invention, and are readily apparent to those with ordinary skill in the art of bioassay.

[0050] The instant invention encompasses the use of the compositions and methods of the instant invention for allergen immunoassays. The hydrophilic PAAEs of the instant invention will be useful for the detection of specific IgE Ab to specific allergens. In this application, hydrophilic PAAEs are linked to streptavidin, and can be used to detect specific allergen-biotin molecules bound to specific antibodies to the allergen. (See an example for the architecture of an ACS specific IgE assay, Fig. 9.)

[0051] Thus the instant invention encompasses methods of measuring the amount of analyte comprising a sample and detecting an analyte specifically and in proportion to its concentration in a sample, and measuring a signal which is generated by a hydrophilic acridinium ester chemiluminescent label that is directly or indirectly proportional to the concentration of the analyte in a sample.

[0052] Such methods of detecting an analyte in a sample comprise contacting a sample with a hydrophilic acridinium ester labeled detector molecule, sequestering bound detector and analyte, washing away excess detector, and measuring a signal from an analyte bound detector.

[0053] Further, methods of detecting an analyte in a sample comprising, contacting the sample with a hydrophilic acridinium ester labeled competitive tracer, and a specific binder for an analyte, retrieving the specific binder, and determining a signal generated by the bound tracer. (See an example for the assay architecture of an ACS testosterone assay, Fig. 7.) Also encompassed are methods of detecting an analyte in a sample comprising contacting a sample with a hydrophilic acridinium ester labeled competitive tracer, and a specific binder for an analyte, retrieving the specific binder, and determining a signal generated by the unbound tracer.

[0054] Included are methods of detecting an analyte in a sample comprising, contacting a sample with a first specific binder, and a second hydrophilic acridinium ester labeled specific binder, and detecting a signal. These assay architectures can be either in the form of sandwich assays or assays in which the second binder (or antibody) is reactive with the first binder. Many variations of these architectures will be apparent to those with ordinary skill in the art.

[0055] The instant invention encompasses methods of detecting an analyte in a sample comprising, contacting a sample with hydrophilic acridinium ester labelled detector and a competitive binder to the detector, sequestering the detector bound to the competitive binder, washing away excess detector and the detector bound analyte, and measuring a signal from the detector bound to the competitive binder. (See an example for the assay architecture of an ACS

estradiol assay, Fig. 8.)

[0056] The methods of the instant invention can be practiced with or without the addition of releasing agents prior to detection of a signal, or other agents which will enhance the signal generated.

[0057] A further aspect of the instant invention encompasses the use of the compositions and methods of the instant invention for gene probe assays. For example, low level concentration of target nucleic acids (the analyte) can be detected by hybridizing to a first nucleic acid probe, which is conjugated with a macromolecular carrier, such as bovine serum albumin, glycoproteins, etc., labeled with multiple hydrophilic PAAEs and optionally multiple hydrophilic polymers such as polyethylene glycol (PEG), and capturing with a second nucleic acid probe, which is immobilized on a solid phase such as paramagnetic particles. The two nucleic acid probes can be preferentially ligated with ligase after the hybridization step, but prior to the separation step (which removes the unbound probe tracer remaining in the supernatant) to enhance assay specificity. Because of the greater number of PAAE incorporated in the tracer due to the hydrophilic nature, good solubility is maintained. Such nucleic acid detection can be accomplished using several schemes that one with ordinary skill in the art can conceive, and using the teachings of the instant invention, develop the proper hydrophilic PAAE tracers for sensitive detection.

[0058] The following examples are presented to illustrate the present invention.

EXAMPLE 1A

Improved synthesis of 2',6'-Dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 9-acridinecarboxylate (DMAeE-NHS, Precursor of DMAE-NHS and NSP-DMAE-NHS)

Succinimidyl 3,5-dimethyl-4-hydroxybenzoate

[0059] A solution of 3,5-dimethyl-4-hydroxybenzoic acid (12.0 g, 72.21 mmol) in 400 mL of anhydrous tetrahydrofuran and 200 mL of anhydrous N,N-dimethylformamide was cooled in an ice-water bath and treated with N-hydroxysuccinimide (8.31 g, 72.21 mmol) with stirring for 15 minutes, followed by addition of dicyclohexylcarbodiimide (17.88 g; 86.65 mmol). After 16 hours of stirring at 25°C under nitrogen, the mixture was filtered; and the wet cake was washed with 10 mL of anhydrous N,N-dimethylformamide. The combined filtrate was evaporated to dryness under reduced pressure. The residue, after washing with anhydrous diethyl ether (2 x 50 mL), was suspended in 80 mL of boiling ethyl acetate for 5 minutes. Cooling the suspension to 25°C gave 11.95 g (63% yield) of the off-white product, which was homogeneous on TLC (R_f 0.6; silica gel, ether).

9-Acridinecarbonyl chloride hydrochloride

[0060] A mixture of 9-acridinecarboxylic acid hydrate (10.0 g, 44.80 mmol) in 80 mL of thionyl chloride was refluxed at 100°C under nitrogen for 2 hours, and then cooled to 25°C. The resulting solution was reduced to about half the original volume under reduced pressure, and then poured into 500 mL of anhydrous diethyl ether. The yellow precipitate was collected, washed with ether (3 x 100 mL), and dried under reduced pressure for 2 hours, to produce 12.95 g (-100% yield) of the product, which was used immediately for the next reaction.

2',6'-Dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 9-acridinecarboxylate (DMAeE-NHS)

[0061] A solution of succinimidyl 3,5-dimethyl-4-hydroxybenzoate (11.29 g, 42.88 mmol) and 4-dimethylaminopyridine (2.19 g, 17.93 mmol) in 200 mL of anhydrous pyridine was cooled in an ice-water bath, followed by addition of 9-acridinecarbonyl chloride hydrochloride (12.95 g, 44.80 mmol). The solution was heated with stirring under nitrogen at 100°C for 2 hours, and then at 25°C for 16 hours. After removal of the solvent under reduced pressure, the residue was flash-chromatographed on a silica gel column (Baker silica gel, Cat#7024-1; F7 x 50 cm) packed with 50% diethyl ether/hexane, and eluted with 75% diethyl ether/hexane (2.5 L), diethyl ether (2 L), 5% ethyl acetate/diethyl ether (3 L) and 25% ethyl acetate/methylene chloride (5 L). Fractions containing the product (R_f 0.4; silica gel TLC plate, ether), collected from the 25% ethyl acetate/methylene chloride eluent, were combined, and evaporated to dryness under reduced pressure, to collect the light-yellow product (DMAeE-NHS) in 10.03 g (50% yield from 9-acridinecarboxylic acid hydrate).

EXAMPLE 1B**Improved synthesis of 2',6'-Dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 10-Methyl-9-acridiniumcarboxylate Methosulfate (DMAE-NHS), Using Toluene as Solvent**

[0062] This procedure uses toluene as a solvent in methylation of DMAeE-NHS, to replace the ozone-depleting reagent 1,1,1-trichloroethane, which was previously used as the solvent in the methylation process as described in U. S. Patent #4,745,181.

[0063] A suspension of DMAeE-NHS (1.0 g, 2.13 mmol) in 50 mL of anhydrous toluene was heated to 110°C to give a homogeneous solution, which was cooled to room temperature followed by addition of 11.0 mL (116.25 mmol) of distilled dimethyl sulfate. After 20 hours of stirring at 110°C under nitrogen, the solution was cooled to room temperature, and then at 4°C for 1 hour. The mixture was filtered, and the yellow wet cake was washed with toluene (2 x 5 mL) followed by anhydrous methylene chloride, heated to boiling for 2-3 minutes, and filtered. The volume of filtrate was reduced to about 50 mL on a hot plate. Cooling the concentrate to room temperature gave yellow crystals, which were collected and washed with diethyl ether (3 x 20 mL), to afford 500 mg (39%) of the pure product. FAB/MS (483, M+).

EXAMPLE 2**Synthesis of 2',6'-Dimethyl-4'-(N-succinimidylloxycarbonyl)phenyl 10-(3'-sulfopropyl)-acridinium-9-carboxylate (NSP-DMAE-NHS)**

[0064] DMAeE-NHS (500 mg, 1.07 mmol) and 1,3-propane sultone (6.5 g, 53.3 mmol) were heated at 150°C under nitrogen in a sealed tube for 20 h. After cooling, excess 1,3-propane sultone was removed by trituration with toluene (3 x 5 mL). The crude product was purified by RP-HPLC using a C18, 30 x 500 mm, 10 µm, preparative column, and eluted with a mixture of 65% solvent A and 35% solvent B. Conditions: solvent A: 0.05% (v/v) TFA/H₂O, solvent B: 0.05% (v/v) TFA/CH₃CN; flow rate 25 mL/min; UV detection at 260 nm. The product was eluted at 26.8 min; yield: 60 mg (25%); FAB/MS (591, M+H).

EXAMPLE 3**Synthesis of 19-Carboxymethyl-17β,19-dihydroxy-4-androsten-3-one (19-Hydroxy-19-carboxymethyl-testosterone, 19-HCMT)**3,17-Bis(ethylenedioxy)-19-ethoxycarbonylmethyl-19-hydroxy-5-androstene

[0065] A 50 mL 2-neck flask equipped with a rubber septum, magnetic stirring bar and N₂ inlet was charged with lithium bis(trimethylsilyl)amide (12.9 mL of 1.0 M solution, 12.9 mmol). The solution was cooled to -78°C under N₂ in dry ice-acetone bath and anhydrous ethyl acetate (1.2 mL, 12.9 mmol) was added dropwise from a syringe. The reaction was stirred at -78°C for 30 minutes and then a solution of 3,17-bis(ethylenedioxy)-5-androsten-19-al (1g, 2.58 mmol) (Lovett, J.A. et al., 1984, J. Med. Chem. 27:734-740) in anhydrous tetrahydrofuran (15 mL) was added dropwise to the solution of the enolate. After completion of the addition, the reaction was stirred for an additional 1.5 hours in the dry ice-acetone bath under nitrogen. The reaction was then quenched with the addition of saturated ammonium chloride solution (-30 mL) and the resulting suspension was extracted three times with ethyl acetate (3 x 25 mL). The combined ethyl acetate extract was washed once with brine (-30 mL) and was then dried over magnesium sulfate. After filtration, the solvent was removed under reduced pressure to yield an oil which was used as such in the next reaction: yield 1.67 g (crude); mass spectrum (chemical ionization) m/z 477 (M+ H+).

3,17-Dioxo-19-ethoxycarbonylmethyl-19-hydroxyl-5-androstene

[0066] The crude bisketal from above (1.67 g) was dissolved in tetrahydrofuran (20 mL) and 3N perchloric acid (10 mL) was added. The reaction was stirred at room temperature for 3 hours and was then diluted with water (40 mL). The resulting suspension was extracted twice with ethyl acetate (2 x 40 mL). The combined ethyl acetate extract was washed once with water (-40 mL) and was then dried over magnesium sulfate. After filtration, the solvent was removed under reduced pressure to afford the crude product which was purified by preparative TLC using 1:1 ethyl acetate/hexanes as eluent (R_f = 0.17): yield 0.46 g (46% two steps) white foam; mass spectrum (chemical ionization) m/z 389 (M + H+).

19-Ethoxycarbonyl-17 β ,19-dihydroxy-4-androsten-3-one

[0067] In a 50 mL round bottom flask equipped with a magnetic stirring bar and a nitrogen inlet, was placed 3,17-dioxo-19-ethoxycarbonylmethyl-19-hydroxy-5-androsten (50 mg, 0.129 mmol) and anhydrous methanol (5 mL). The solution of the steroid was stirred and cooled in ice under nitrogen and sodium borohydride (6.4 mg, 0.168 mmol) was added as a solution in methanol (1 mL). The reaction was stirred in ice for 8 minutes and was then quenched with the addition of acetone (1 mL) and acetic acid (2 mL). The reaction mixture was then diluted with water (-25 mL) and the resulting suspension was extracted with ethyl acetate (-35 mL). The ethyl acetate extract was dried over magnesium sulfate, filtered and then concentrated under reduced pressure to afford a white powder: yield 51 mg (quant.) which was homogenous on TLC (1:19, MeOH/CHCl₃) (R_f = 0.26). This material was used as such in the next reaction.

19-carboxymethyl-17 β ,19-dihydroxy-4-androsten-3-one (19-HCMT)

[0068] The ethyl ester from above (51 mg, 0.129 mmol) was dissolved in methanol (5 mL) and treated with KOH (0.25 g). The reaction was stirred at room temperature for 1 hour and was then quenched with the addition of water (25 mL) and acetic acid until the pH of the solution was adjusted to 6. The resulting mixture was further acidified with the addition of 0.5 N HCl (1 mL) and the resulting suspension was extracted three times with ethyl acetate (3 x 15 mL). The combined ethyl acetate extract was dried over magnesium sulfate, filtered and concentrated. The crude acid was purified by preparative TLC using 2% acetic acid in ethyl acetate as eluent (R_f = 0.4) and was isolated as a white powder. yield: 10 mg (22%); mass spectrum (chemical ionization) 363 (M + H⁺).

Example 4**Synthesis of 10-Carboxyvinyl-17 β -hydroxy-19-nor-4-androsten-3one (19-Carboxyvinyl-nor-tastosterone; 19-CVNT)**3,17-Bis(ethylenedioxy)-10-ethoxycarbonylviny-19-nor-5-androstene

[0069] A 50 mL 2-neck flask equipped with a magnetic stirring bar, reflux condenser with a nitrogen inlet and a rubber septum was charged with triethylphosphonoacetate (0.35 g, 1.7 mmol) and anhydrous tetrahydrofuran (5 mL). The solution was cooled in an ice-salt bath to -10°C under nitrogen and n-butyl lithium (1.24 mL of 1.6 M solution, 1.7 mmol) was added. The reaction was stirred at -10°C for 30 minutes and then a solution of 3,17-bis(ethylenedioxy)-5-androsten-19-al (0.22 g, 0.57 mmol) in anhydrous tetrahydrofuran (5 mL) was added dropwise from a syringe. The reaction was then warmed to room temperature and was then refluxed under nitrogen for 24 hours. After cooling to room temperature, the reaction mixture was then partitioned between a 1:1 mixture of water and ethyl acetate. The ethyl acetate layer was separated and washed with brine. It was then dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by preparative TLC using 2:3, ethyl acetate/hexanes as eluent (R_f = 0.34) and was isolated as a white fluffy solid: yield 0.11 g (43%); mass spectrum (chemical ionization) 459 (M + H⁺).

3,17-Dioxo-10-ethoxycarbonylviny-19-nor-4-androstene

[0070] A solution of the bisketal (0.11 g) from above in acetone (10 mL) was treated with p-toluenesulfonic acid (20 mg). The reaction was stirred at room temperature for 16 hours and was then concentrated under reduced pressure. The resulting residue was partitioned between ethyl acetate and water. The ethyl acetate layer was separated, dried over magnesium sulfate and then concentrated under reduced pressure. The product was purified by preparative TLC using 1:1, ethyl acetate/hexanes as eluent (R_f = 0.51) and was isolated as a white powder yield 20 mg (23%). This material was used as such for the next reaction.

10-Ethoxycarbonylviny-17 β -hydroxy-19-nor-4-androsten-3-one

[0071] A solution of 3,17-dioxo-10-ethoxycarbonylviny-19-nor-4-androstene (19.5 mg, 0.053 mmol) in methanol (3 mL) was cooled in ice under nitrogen and was treated with sodium borohydride (2.7 mg). The reaction was stirred in ice under nitrogen for 30 minutes and was then quenched with the addition of acetone (1 mL) and acetic acid (1 mL). The resulting solution was concentrated and the residue was partitioned between ethyl acetate and water. The ethyl acetate layer was separated, dried over magnesium sulfate, filtered and then concentrated under reduced pressure. The residue obtained (19 ng) was dissolved in anhydrous chloroform (5 mL) and treated with activated MnO₂ (100 mg). The black suspension was stirred at room temperature under nitrogen 30 minutes and was then diluted with ethyl

acetate and filtered. The filtrate was concentrated under reduced pressure to afford a white powder which was homogenous on TLC (1:19, methanol/chloroform, $R_f = 0.28$): yield 18.2 mg (92%).

10-Carboxyvinyl-17 β -hydroxy-19-nor-4-androsten-3-one (19-Carboxyvinyl-nor-testosterone; 19-CVNT)

[0072] A solution of 10-ethoxycarbonylvinyl-17 β -hydroxy-19-nor-4-androsten-3-one (17 mg) in methanol (3 mL) was treated with 5% KOH (2 mL). The reaction was stirred at room temperature for 30 minutes and was then acidified with 1N HCl. The resulting suspension was extracted twice with ethyl acetate (2 x 25 mL). The combined ethyl acetate extract was dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford a white powder which was homogenous on TLC (2% acetic acid in ethyl acetate, $R_f = 0.25$): yield 15.2 mg (97%); mass spectrum (chemical ionization) 345 (M + H⁺).

Example 5

Synthesis of NSP-DMAE / 19-HCMT Conjugate Cross-linked with n-Hexyl-1,6-diamine (NSP-DMAE-HD-19-HCMT)

[0073] First, NSP-DMAE-NHS was crosslinked with n-hexyl-1,6-diamine to yield NSP-DMAE-HD. Thus, to a yellow solution of NSP-DMAE-NHS (60 mg, 0.1 mmol) in anhydrous DMF / CH₃OH (1:4, 10 mL) was added n-hexyl-1,6-diamine (116.2 mg, 1 mmol). The resulting colorless solution was stirred at room temperature under nitrogen for 15 hours, and concentrated to ~1 mL. The crude product was purified by TLC using a preparative silica gel plate (2000 mm, 20 x 20 cm) developed with CHCl₃ / CH₃OH / H₂O (55 / 40 / 5). The product had $R_f = 0.5$; yield: 50.2 mg (84%); FAB / MS (593, M+H).

[0074] Subsequently, a solution of 19-hydroxy-19-carboxymethyl-testosterone (19-HCMT, 4 mg, 0.011 mmol) in anhydrous DMF (0.1 mL) was diluted with anhydrous CHCl₃ (0.4 mL), cooled in an ice-bath, and activated with a solution of DCC (2.7 mg, 0.013 mmol) in anhydrous CHCl₃ (0.1 mL). The reaction mixture was stirred at 0°C for 10 minutes, followed by the addition of NSP-DMAE-HD (6 mg, 0.01 mmol) in anhydrous DMF (0.2 mL) and the new reaction mixture was stirred at room temperature for 15 hours. The crude product was purified by RP-HPLC using a C18, 7.8 x 300 mm, 10 μ m, semipreparative column, solvent A: 0.05% (v/v) TFA/H₂O, solvent B: 0.05% (v/v) TFA/CH₃CN, flow rate: 2.5 mL/min, UV detection at 260 nm. Initially 30% solvent B was used for 15 minute followed by a linear gradient to 50% solvent B in 5 min. The product was eluted at 24.5 minutes; yield: 2.3 mg (25%); FAB / MS (936, M+H).

Example 6

Synthesis of NSP-DMAE / 19-CVNT Conjugate Cross-linked with n-Hexyl-1,6-diamine (NSP-DMAE-HD-19-CVNT)

[0075] NSP-DMAE-HD-19-CVNT was prepared from 19-CVNT (2 mg, 0.006 mmol) and NSP-DMAE-HD (3 mg, 0.005 mmol) as described in example 5. The crude product was purified by RP-HPLC using a C18, 10 μ m, 7.8 x 300mm semipreparative column. Initially 30% solvent B was used for 15 minutes followed by a linear gradient to 60% solvent B in 5 minutes, where solvent A: 0.05% (v/v) TFA/H₂O and solvent B: 0.05% (v/v) TFA/CH₃CN; flow rate 2.5 mL / min; UV detection at 260 nm. The product was eluted at 23.2 minutes. Product yield: 0.72 mg (16%); FAB / MS (918, M+H).

Example 7

Synthesis of NSP-DMAE / Testosterone-19-CME Conjugate cross-linked with n-Hexyl-1,6-diamine (NSP-DMAE-HD-19-CMET)

[0076] NSP-DMAE-HD-19-CMET was prepared from testosterone-19-CME (4 mg, 0.011 mmol) and NSP-DMAE-HD (3 mg, 0.005 mmol) as described in example 5. The crude reaction mixture was first separated on a preparative TLC plate (silica gel, 1000 μ m, 20% MeOH/CH₂Cl₂). Two bands at R_f 0.2 and R_f 0.3 were collected, combined and further purified by RP-HPLC using a C18, 10 μ m, 7.8 x 300mm semipreparative column. Initially 30% solvent B was used for 15 minutes followed by a linear gradient to 60% solvent B in 5 minutes, where solvent A: 0.05% (v/v) TFA/H₂O and solvent B: 0.05% (v/v) TFA/CH₃CN; flow rate 2.5 mL / min; UV detection at 260 nm. The product was eluted at 24.1 minutes. Product yield: 0.3 mg (6.4 %); FAB / MS (936, M+H).

EXAMPLE 8**Synthesis of DMAE/19-HCMT conjugate cross-linked with n-hexyl-1,6-diamine (DMAE-HD-19-HCMT)**

[0077] First, DMAE-HHS was derivatized with n-hexyl-1,6-diamine to yield DMAE-HD as described in Example 5. Thus to a yellow solution of DMAE-NHS (so mg, 0.084 mmol) in anhydrous DMF/CH₃OH (1:4, 10 mL) was added n-hexyl-1,6-diamine (98 mg, 0.84 mmol). The resulting solution was stirred at room temperature under nitrogen for about 4 hours and evaporated. The crude product was purified by TLC using preparative silica gel plates (2000 mm, 20 x 20 cm) developed with CHCl₃/CH₃OH/H₂O (65 / 25 / 4). The product band at R_f = 0.4 was stripped, eluted with the same solvent system. The eluent was evaporated to a residue which was triturated with chloroform, filtered and the filtrate evaporated to give purified DMAE-HD (20 mg, 41%).

[0078] DMAE-HD-19-HCMT was prepared from 19-HCMT (4.75 mg, 0.013 mmol) and DMAE-HD (7 mg, 0.012 mmol) as described in Example 5. One third of the crude product was purified by RP-HPLC using a C18, 10 μ , 7.8 x 300 mm semipreparative column. Initially 30% solvent B was used for 15 minutes followed by a linear gradient to 60% solvent B in 5 minutes, where solvent A: 0.1% (v/v) TFA/H₂O and solvent B: 0.1% (v/v) TFA/CH₃CN, flow rate 2.3 mL/min; UV detection at 260 nm. The product was eluted at 23 min. Product yield: 1.5 mg (45 %); FAB/MS (828.5 M⁺).

EXAMPLE 9**Assay For Testosterone**

[0079] This example teaches a sensitive and rapid competitive chemiluminescence immunoassay for testosterone using the hydrophilic NSP-DMAE-HD-19-HCMT tracer, that is heterologous to the antibody immunogen at the bridge. The data shows the clear advantage of this new hydrophilic tracer compared to the conventional hydrophobic DMAE-HD-19-HCMT tracer by allowing the development of a rapid, high sensitivity testosterone assay.

A) Production and characterization of polyclonal and monoclonal antibodies:

[0080] commercially available, rabbit polyclonal antibodies that are anti-testosterone-19-o-carboxymethylether-Human Serum Albumin (RAT-19), was purchased from OEM, (Tom's River, NJ). The production of this antibody was previously described by Rao (U.S. Patent # 4,197,286). A mouse monoclonal antibody against rabbit IgG was produced by a modified version of the technique described by Kohler and Milstein (Nature (London) 256, 495-497 (1975)). Mice were immunized with purified rabbit IgG to produce a mouse monoclonal anti-rabbit IgG (MAR-IgG). Standard antibody screening techniques were used to select a hybridoma with high specificity for rabbit IgG. Ascites was produced using clone HMF1-11B5-3F1-5H7. MAR-IgG antibody was purified from the ascites using Protein A-sepharose (Pharmacia Fine Chemicals, Piscataway, N.J.) affinity chromatography.

B) Development of a Testosterone Chemiluminescence Immunoassay

[0081] Optimal assay performance for a rapid testosterone chemiluminescence assay was achieved with the MAR-IgG (described above) covalently coupled to a solid phase and preincubated with the RAT-19 antibody using the NSP-DMAE-HD-19-HCMT as the tracer. The assay used paramagnetic particles (PMP) (Advanced Magnetics Inc., Cambridge, MA) as the solid phase.

[0082] The MAR-IgG (160 mg per gram of PMP) in 0.1 M sodium phosphate, pH 7.4, was covalently linked to the PMP after activation of the PMP with 0.5% glutaraldehyde in 0.1 N sodium phosphate (method modified from Groman *et al.*, 1985, BioTechniques 3, 156-160). This was followed by incubation with RAT-19. Excess RAT-19 was removed by additional washes of the reaction mix with phosphate buffer.

[0083] Twenty-five microliters (mL) of a serum sample or testosterone standard (0.1 to 24 ng/mL or 2.5 to 600 pg/tube), 50 mL of NSP-DMAE-HD-19-HCMT tracer (6 x 10⁶ Relative Light Units/test) and 300 mL of MAR-IgG/RAT-19 antibody-solid phase were added simultaneously, incubated at 37°C for 5 minutes, washed and the chemiluminescence measured on an ACS:180 (Ciba Corning Diagnostics, Medfield, MA) automated chemiluminescence immunoassay system. The chemiluminescence signal (Relative Light Units) is inversely proportional to the testosterone concentration. Assay standards consisted of pure testosterone in steroid-free, charcoal stripped human plasma. The testosterone standard binding curve (percent of RLU of standard to RLU of zero standard) for the described assay is shown in FIG. 3. Testosterone doses from serum samples were generated on the ACS:180 using two point calibration off a master curve. Three other testosterone tracers, namely, NSP-DMAE-HD-19-CVNT (heterologous to the antibody immunogen), NSP-DMAE-HD-19-CMET (homologous to the antibody immunogen), and DMAE-HD-19-HCMT (a non-hydrophilic version of the NSP-DMAE-HD-19-HCMT) were evaluated in the same assay format for their effect on the binding curve

and the 50% maximum binding (ED50) of the standard binding curve FIG. 4. The DMAE-HD-19-HCMT tracer was evaluated to demonstrate the advantage of employing a hydrophilic PAAE (NSP-DMAE) tag versus a hydrophobic PAAE tag (DMAE).

C) Conventional Radioimmunoassay for Testosterone

[0084] The results (n=145) of a conventional testosterone radioimmunoassay (¹²⁵Iodine) were compared in selected male and female serum samples with the ACS:180 chemiluminescence assay using the heterologous testosterone tracer, NSP-DMAE-HD-19-HCMT. The Coat-A-Count Total Testosterone (DPC, Los Angeles, CA) commercial system was used. This assay has a three hour incubation time with a 50 mL sample size. The reported minimum detectable dose is approximately 2 picograms per tube as calculated from the manufacturers package insert V116 dated January 27, 1992. This assay was performed according to the manufacturer's instructions.

D) Competitive Chemiluminescence Assay

[0085] The minimum detectable dose (Rodbard, 1978 Anal. Biochem. 90, 1-12) of the rapid automated testosterone immunoassay using the heterologous testosterone tracer NSP-DMAE-HD-19-HCMT is 1.25 pg/tube, a 1.6 fold increase over the conventional 3 hour radioimmunoassay method described above. As shown in FIG. 5, patient sample results for the chemiluminescent assay correlate very well to the radioimmunoassay. The crossreactivity profile of the assay is shown in Table 2. These data suggest that the new chemiluminescent assay is of useful value in the analytical determination of testosterone in human serum samples.

TABLE 2

Specificity of ACS Testosterone chemiluminescence immunoassay using NSP-DMAE-HD-19-HCMT tracer			
Cross-reactant	Dose	% Cross-reactivity	
Specifications %			
Androstenedione	100 ng/mL	1.55	<1
Androsterone	1 ug/mL	0.03	<0.1
cortisol	1 ug/mL	0.03	<0.1
Corticosterone	1 ug/mL	0.02	<0.1
11-deoxycortisol	1 ug/mL	0.05	<0.1
5α-dihydrotestosterone	100 ng/mL	8.3	<15
Estradiol-17B	100 ng/mL	0.12	<0.1
Estrone	100 ng/mL	0.07	<0.1
Progesterone	1 ug/mL	0.09	<0.1
Danazol	1 ug/mL	0.06	<0.1
Dexamethasone	1 ug/mL	0.01	<0.1

[0086] A comparison of the standard binding curves and calculated ED50's for each chemiluminescent tracer under the same conditions are shown in FIG. 4. The highest ED50 is for the NSP-DMAE-HD-19-CMET tracer that is homologous to the antibody immunogen. The ED50's, for both heterologous tracers, NSP-DMAE-HD-19-CVNT and NSP-DMAE-HD-19-HCMT, are lower than for the homologous tracer indicating that the heterologous tracers are more easily displaced by testosterone than the homologous tracer. Little displacement is seen with the hydrophobic DKAE-HD-19-HCMT tracer indicating the clear advantage of the hydrophilic NSP-DMAE-HD-19-HCMT version of the tracer in testosterone immunoassay.

EXAMPLE 10

RUBELLA IgG CAPTURE ASSAY

[0087] A rubella IgG capture assay was used to evaluate NSP-DMAE-NHS and DMAE-NHS as reagents for the direct labelling of rubella. Intact rubella virus was labeled with either NSP-DMAE-NHS or DMAE-NHS as described below. Both of these tracer preparations were subjected to a titering study and a population study. The performance

of each rubella tracer can be determined by analyzing the positive sample/negative sample ratio generated by each tracer for a given sample population. In this rubella IgG capture assay, the Positive signal/Negative signal (P/N) ratio also reflects on the ability to accurately distinguish small differences in the level of rubella specific IgG. The preferred rubella tracer is one that generates appreciable differences in P/N ratios for samples spanning a wide range of IU/mL levels, since this will lead to greater sensitivity in rubella specific IgG detection.

A) Preparation of Rubella virus Tracers

[0088] Rubella virus (grade IV, 1 mL at 0.5 mg/mL) purchased from Viral Antigens, Inc (Memphis, TN) was dialyzed into labelling buffer (10 mM Phosphate, 0.125 N NaCl, pH 8) using a centricon-30 microconcentrator (Amicon, Danvers, MA). The virus sample was brought to 2 mL volume in labelling buffer, then concentrated 10-fold by spinning the centricon-30 in a tabletop centrifuge for 30 minutes at 2600 rpm. The concentrate was again brought to 2 mL with labelling buffer. This concentration and reconstitution process was repeated 3 times. The sample was then split into . 2 x 1 mL aliquots for use in subsequent labelling procedures.

[0089] To each Rubella virus aliquot (0.25 mg in 1 mL buffer) was added NSP-DMAE-NHS (50 µg in 25 µL dimethylformamide) or DMAE-NHS (50 µg in 25 µL dimethylformamide) solution separately. The labelling reaction mix was vortexed briefly and incubated at room temperature for 15 minutes and stopped by adding 5 mg of lysine (Sigma, St. Louis, MO) in 50 µL of water. The reaction mixture was vortexed briefly, incubated at room temperature for another 5 minutes and subjected to following purification steps.

[0090] The reaction mixture was first purified through PD10 Sephadex G25M sizing column (Pharmacia, Piscataway, NJ) preequilibrated and eluted with labelling buffer containing 0.1% BSA. The flow rate of the elution was maintained at 1 mL/min and 20 x 0.8 minute fractions were collected. The tracer which appeared in the void volume as the first RLU peak (fractions 6-9) was located by taking a 10 µL sample from each fraction, diluting it 200-fold with labelling buffer, and counting 10 µL of the diluted sample on a MLA II luminometer (Ciba Corning Diagnostics, Medfield, MA). The desired pooled fractions were concentrated with a centricon-30 to about 300 µL. The concentrate was subjected to further HPLC gel filtration.

[0091] Waters HPLC system (Waters Associate, Milford, MA) mounted with a Protein Pak 300SW HPLC gel filtration column (Waters #11787) was first equilibrated with HPLC column buffer containing 50 mM Tris pH 8, 0.25 M MgCl₂, 0.1 % BSA, and 0.01% Triton X-100. The labeled virus concentrate from the PD10 column was loaded onto the 300SW column. The elution was maintained at a flow rate of 0.5 mL/min and 40 x 0.25 mL fractions were collected beginning from 11 minute after sample injection. The labeled virus peak (retention times of 13-17 minutes) was again identified in the same manner as above, by looking for the first RLU peak through counting samples of the collected fractions. The fractions containing the first RLU peak were pooled and represented the purified labeled virus tracer.

B) Preparation of Anti-human IgG solid phase

[0092] Mouse anti-human IgGFc monoclonal antibody was covalently linked to PMP with glutaraldehyde (in 20 mM sodium phosphate, pH 7). This is a modification of the method described in BioTechniques 3, 156 (1985).

c) Preparation of monoclonal anti-Human IgGFc (clone 157.8D6.5D3)

[0093] Fusion Procedure: Method uses polyethylene glycol to fuse isolated mouse spleen cells from immunized donors with SP2/0 Ag4-1 mouse myeloma cells (Galfrie and Milstein, (1981), *Methods in Enzymology* 73:3).

[0094] Screening Procedure: Secretion of specific antibody was tested for by the ability of the cell supernatant to link acridinium ester labeled human IgG to human IgG immobilized on paramagnetic particles. The monoclonal antibody was shown to be specific for human IgGFc and nonreactive to human IgG Fab and Fab₂ fragments, human IgA, IgM, IgE, and IgD when tested by western blot (as per Proc. Nat. Acad. Sci. U.S.A., 76: 4350 (1979)).

[0095] Antibody Isotype: The antibody isotype is IgG1, identified by using the Zymed MonoAb ID EIA kit (Zymed Lab Inc., San Francisco, CA).

D) Preparation of monoclonal Anti-Rubella antibody (clone ML109.10E7.4A3)

Fusion Procedure: same as above.

[0096] Screening Procedure: Secretion of specific antibody was tested for by the ability of the cell supernatant to bind rubella virus. The monoclonal binding to antigen was detected by ELISA (as per J.E. Coligan, *Current Protocols in Immunology Vol. 1* (1991) section 2.1.3). Clones were shown to be specific by nonreactivity to uninfected culture controls.

E) Rubella IgG Capture Chemiluminescence Immunoassay

[0097] Performance of the different rubella virus tracers was evaluated in a manual Magic Lite® (Ciba Corning Diagnostics, Medfield, MA) assay format in triplicate. All serum samples were diluted five-fold with the diluent (1 x DPBS, BioWhittaker, Walkersville, MD) containing 50% calf serum, 1% Triton X-100, 1% normal mouse serum, and 0.1% sodium azide. Diluted sample (50 µL), Anti-human IgG solid phase (80 µg in 250 µL), and diluted rubella virus tracer (100 µL) were incubated together for 7.5 minutes at 37°C in a Sarstedt polystyrene tube (Newton, NC). The reaction mix was separated, aspirated and washed twice with 1 mL 1 x DPBS (BioWhittaker, Walkersville, MD). The final pellet was reconstituted with 100 µL of water and its chemiluminescence measured on MLA-II luminometer (Ciba Corning Diagnostics, Medfield, MA). Since the two rubella virus tracers (prepared as described above) varied in their specific chemiluminescence activity, the titering study was conducted over different ranges of input counts, using nine rubella IgG samples as shown in Table 3.

TABLE 3

Rubella IgG Capture Assay Results of NSP-DMAE Rubella Titering Study						
Sample	NSP-DMAE Rubella (RLU, %CV, P/N)					
	1x10 ⁶	5.95x10 ⁶	11.1x10 ⁶	13.3x10 ⁶	24.3x10 ⁶	46.1x10 ⁶
<i>Diluent</i>	1233	1793	2473	2757	4253	9360
%CV	1.9	2.6	9.4	3.3	5.5	17.2
P/N	1.0	0.9	1.0	0.8	1.1	1.6
<i>Negative</i>						
RAB 17	1233	2037	2447	3367	4023	5987
%CV	3.3	20.1	3.9	23	8.5	7.5
P/N	1.0	1.0	1.0	1.0	1.0	1.0
RAB 36	1383	1897	2587	2880	6357	8357
%CV	13.3	13.2	4.7	0.9	16.3	21.7
P/N	1.1	0.9	1.1	0.9	1.6	1.4
<i>Low +</i>						
CAP 10	1393	2903	4247	5613	8100	14830
%CV	2.7	6.1	2.4	6.4	9.1	7.6
P/N	1.1	1.4	2.2	2.1	2.6	3.3
CAP 15	1473	3170	5283	7003	10263	19737
%CV	2.7	3	3.2	2.7	6.3	7.9
P/N	1.2	1.6	2.2	2.1	2.6	3.3
<i>Mid-High+</i>						
TSI 3156	3547	14420	32143	40403	66857	142657
%CV	3.8	3.3	4	3.2	1.3	1
P/N	2.9	7.1	13.1	12.0	16.6	23.8
RAB 67	10970	52003	120987	151000	262317	536050
%CV	2.7	6.4	5	2.4	1.06	2.3
P/N	8.9	25.5	49.4	44.8	65.2	89.5
TSI 3167	7357	35337	75027	94530	167653	342447
%CV	1.9	1.9	8	3.4	4.6	3.5
P/N	6.0	17.3	30.7	28.1	41.7	57.2
TSI 3163	11270	56443	111327	142290	249450	514440
%CV	2.2	3.3	4.9	1.7	2.2	10.6

EP 0 754 178 B1

TABLE 3 (continued)

Rubella IgG Capture Assay Results of NSP-DMAE Rubella Titering Study						
Sample	NSP-DMAE Rubella (RLU, %CV, P/N)					
<i>Mid-High+</i>						
P/N	9.1	27.7	45.5	42.3	62.0	85.9
DL1-3150-388	11477	55093	113783	146473	263267	542483
%CV	1.4	6.5	12.3	3.2	3.2	3.4
P/N	9.3	27.0	46.5	43.5	65.4	90.6

[0098] The DMAE labeled rubella tracer was titered over a wider range due to a 4-5 fold greater degree of incorporation of DMAE into the labeled virus tracer compared to NSP-DMAE. In order to compare assay conditions using a comparable mass of rubella tracers, the DMAE rubella tracer total input range must be several times higher than that of NSP-DMAE rubella condition. The titering study results demonstrated improved P/N ratios as the level of NSP-DMAE rubella tracer increased, while the P/N ratios generated by DMAE rubella tracer plateaued at 10×10^6 RLU/test. The selection of 24×10^6 RLU/test as the optimal input level for the NSP-DMAE rubella tracer was based on the results of higher P/N ratios, while maintaining low nonspecific binding with the control diluent. The comparable assay conditions using DMAE rubella tracer is 125×10^6 RLU/test. When these two conditions are compared, it is clear that NSP-DMAE rubella tracer is the superior reagent with P/N ratios ranging from 2 to 4 fold greater than those generated by DMAE rubella tracer.

[0099] The testing of these two rubella tracers was expanded to a population of twenty rubella IgG samples (ranging from 0.3-500 IU/mL) and 4 potential crossreactive samples (all identified as rubella IgG negative by Rubazyme assay (Abbott Labs, Chicago, IL) (Table 4A).

The study was conducted using the following total inputs for each reagent; NSP-DMAE rubella tracer (24×10^6 RLU/test) and DMAE rubella tracer (125×10^6 RLU/test). The population data suggested that the sensitivity of the rubella IgG capture assay is affected markedly by the class of the PAEE utilized to directly label the virus. The P/N (positive to negative result) ratios generated from employing NSP-DMAE rubella tracer are considerably greater than those generated by the DMAE rubella tracer (FIG. 6). These results reflect the ability of the assay performed with hydrophilic NSP-DMAE rubella tracer to more clearly distinguish positive samples from negative samples than conventional DMAE tracer. This is particularly important in the diagnostically critical low positive population.

[0100] The impact of hydrophilic NSP-DMAE rubella tracer on the specificity of the potentially crossreactive samples was evaluated. Using a P/N ratio cutoff corresponding to the CAP10 sample, the results indicate no false positive identifications resulting from the use of the rubella tracers with these types of samples (Table 4B).

[0101] The results of both of these studies clearly suggest NSP-DMAE-NHS is a superior reagent for the direct labelling of rubella virus. This reagent results in a rubella preparation which specifically binds to a wide range of rubella IgG levels to produce specific RLU values which are clearly distinguishable from those of a rubella IgG negative population. Furthermore, these conditions exist without incurring false positive results with potentially crossreactive samples.

TABLE 4A

Population study using NSP-DMAE Rubella and DMAE Rubella Tracers		
Sample	Rubella IgG Capture Assay (P/N ratio)	
	NSP-DMAE Rubella	DMAE Rubella
RAB17	1.0	1.1
AP05874-4	1.0	1.0
AS04912-9	1.1	1.1
RAB 36	1.1	1.2
AS14837-8	1.3	1.0
CAP 10	1.8	1.2
AS01721-3	2.1	1.0
AP27771-3	2.4	1.0
AR99860-4	2.5	0.8

EP 0 754 178 B1

TABLE 4A (continued)

Population study using NSP-DMAE Rubella and DMAE Rubella Tracers		
Sample	Rubella IgG Capture Assay (P/N ratio)	
	NSP-DMAE Rubella	DMAE Rubella
CAP 15	2.5	1.0
TSI 3164	5.8	1.8
AS06137-4	8.0	1.3
GP1-27	23.9	2.8
AP41719-0	24.9	3.5
GP2-9	26.2	3.1
GP2-18	27.7	3.4
AR86731-7	51.1	5.6
AR96863-2	58.7	5.9
AS15758-2	68.5	6.9
AR74744-4	88.9	9.8

TABLE 4B

Interference Samples Rubazyme (Index) (All Rubella IgG -)	Rubella Assay (P/N ratio)		
	NSP-DMAE	DMAE	>= 1.0 is +
EBV+E9-3204-467	1.5	1.1	0.725
EBV+E8-2405-99	1.01	0.98	0.171
VZV+E02606-135	1.4	1.1	0.183
ANA+12615M-910513	1.7	1.1	0.793

F) Comparison with Ciba Corning Magic Lite® control Assay and Abbott IMx Assay

[0102] The evaluation of NSP-DMAE rubella tracer was extended even further, to include an investigation of how this Magic Lite® rubella IgG assay using NSP-DMAE rubella tracer compares with (1) Ciba Corning Diagnostics Magic Lite® assay (which uses a complex of unlabelled rubella and NSP-DMAE labeled anti-rubella monoclonal antibody 10E7) and (2) IMx Rubella IgG (Abbott, Chicago, IL). The sample population tested by both Magic Lite® assays included thirty-nine rubella IgG samples identified as positive or negative for rubella specific IgG by IMx (Table 5). It is interesting to note that the Positive result/Negative result (P/N) ratios generated by NSP-DMAE rubella tracer are significantly greater than observed with the Magic Lite® control assay format. This is particularly evident in the rubella IgG high positive samples and may reflect competition between the rubella specific IgG and the NSP-DMAE-MAb10E7 tracer for binding sites on the unlabelled rubella virus. Based on a P/N ratio cutoff corresponding to the CAP10 sample, the Magic Lite® rubella IgG assay utilizing NSP-DMAE rubella tracer demonstrated 97.4% agreement with the Magic Lite® rubella control assay and 92.3% agreement with IMx rubella IgG.

TABLE 5

Expanded population study using NSP-DMAE Rubella Tracer			
Sample	NSP-DMAE	Rubella +	IMx
	Rubella	NSP-DMAE10E7	
RAB 17	1.0	1.0	negative
AP05874-4	1.0	0.9	negative
AS04912-9	1.1	0.9	negative
RAB 36	1.1	1.0	negative
AS14837-8	1.3	0.9	negative

EP 0 754 178 B1

TABLE 5 (continued)

Expanded population study using NSP-DMAE Rubella Tracer			
Sample	NSP-DMAE	Rubella +	IMx
	Rubella	NSP-DMAE10E7	
CAP 10	1.8	1.1	positive
AS01721-3	2.1	1.05	negative
AP27771-3	2.4	1.1	negative
AR99860-4	2.5	1.1	negatade
CAP 15	2.5	1.2	positive
AS00541-9	2.8	1.2	positive
AR97400-3	3.2	1.3	positive
AP41598-1	4.0	1.5	positive
TSI 3164	5.8	2.0	positive
AS06137-4	8.0	1.7	positive
AS00407-3	9.9	2.4	positive
TSI 3156	13.6	3.0	positive
GP2-3	23.4	3.7	positive
GP1-27	23.9	3.6	positive
AP41719-0	24.9	2.6	positive
GP2-9	26.2	3.9	positive
GP1-1	26.3	3.7	positive
GP2-11	26.9	4.1	positive
GP2-18	27.7	4.1	positive
GP2-4	28.1	4.4	positive
TSI 3167	33.4	6.2	positive
GP2-22	39.7	5.4	positive
GP2-14	42.0	4.1	positive
GP1-17	43.3	7.4	positive
GP2-35	44.6	7.8	positive
TSI 3163	48.7	13.1	positive
AS15788-7	49.3	5.3	positive
DL1-3510-388	51.0	11.6	positive
AR86731-7	51.1	9.6	positive
AR96863-2	58.7	10.6	positive
AP35925-6	62.0	7.9	positive
HG68085-5	63.8	16.5	positive
AS15758-2	68.5	11.3	positive
AR74744-4	88.9	27.9	positive

EXAMPLE 11.

DMAE & NSP-DMAE labeling of protein

[0103] This example shows that a larger quantity of NSP-DMAE can be covalently attached to a protein, such as BSA (bovine serum albumin) because of the hydrophilic nature of NSP-DMAE. A solution of BSA (10 mg, 150 nmol) in 0.1 M carbonate buffer, pH 9, (2.7 mL) was treated with a solution of 50 equivalents of either DMAE-NHS ester or NSP-DMAE-NHS ester (4.5 mg each, 7.5 mmol) in DMF (300 mL). The reaction utilizing DMAE-NHS was slightly cloudy whereas the reaction with NSP-DMAE-NHS remained completely clear throughout the reaction. The reactions were stirred at room temperature for 24 hours and the labeled proteins were isolated by size-exclusion chromatography on a Sephadex G-25 column (3.7 x 42 cm) eluted with 10 mM phosphate buffer pH 8. The protein fraction was collected in each case, concentrated and repurified on the same Sephadex G-25 column. After the second concentration, the quantity of the labeled protein recovered from each reaction and the extent of Acridinium Ester (AE) labeling was

estimated by performing a protein analysis (Bio-Rad protein assay) and by measuring the chemiluminescent activity of the labeled protein, respectively. From these measurements, the protein was found to be labeled with approximately 14 DMAE molecules and 24 NSP-DMAE molecules.

Example 12

Conjugation of NSP-DMAE labeled BSA (Bovine Serum Albumin) to 5'-32P-labeled 508.CF-3'-Maleimide:

[0104] This example shows how the protein carrier pre-labeled with plurality of NSP-DMAE can be covalently attached with a 24-mer oligonucleotide which we called 508.CF-NH₂. The oligonucleotide was first radio-labeled with 32-P at the 5'-end phosphate using the conventional radio-labeling technique commonly known to those skilled in the arts. The purpose of 32P-labeling of the oligonucleotide was to provide a handle to quantify the oligonucleotide once conjugated with the protein carrier. The 3'-end of the oligonucleotide carries an aminoalkyl group to allow the reaction with a bi-functional crosslinker as described below so that a sulfhydryl-reactive maleimido group can be attached. Methods of preparing oligonucleotide with aminoalkyl group at the 3'-end is well known (See User's Manual on DNA Modification Reagents for Use in Automated DNA Synthesis, Doc. No. PB022789-1, CLONTECH Laboratories, Inc., Palo Alto, CA, 1989).

Preparation of 32P-labeled 508.CF-maleimide:

[0105] A solution of 32P-508.CF-NH₂ (40 nmoles) in 50 mM carbonate. (600 mL, pH 8.4) was treated with sulfo-succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (sulfo-SMCC, 10 mg, 22.9 mmol). The reaction was stirred at room temperature for 30 minutes following which the product was purified directly by preparative HPLC on a C8 column (0.7 x 25 cm) using a gradient of 8% to 20% acetonitrile in 0.1 M triethylammonium acetate buffer, pH 7.0, over 20 minutes at a flow of 2.3 mL/min and UV detection at 260 nm (t_R = 14.8 min. product; t_R = 12.7 min. starting material). The eluent off the HPLC, containing the product was lyophilized to dryness: yield 25 nmoles (60%).

Conjugation of the NSP-DMAE labeled BSA to 32P-508.CF-maleimide:

[0106] (NSP-DMAE)₂₁-BSA was prepared by reacting BSA with 30 equivalents of NSP-DMAE-NHS as described above. The labeled protein (6mg, 90 nmole) in 100 mM phosphate buffer containing 5 mM EDTA, pH 8, was treated with 2-iminothiolane (50 equivalents, 0.6 mg, Pierce, Rockford, IL).

[0107] The reaction was stirred under nitrogen at room temperature for 1 hour. The thiolated protein was then isolated by size-exclusion chromatography on a column of Sephadex G-25 (3.7 x 42 cm) using 20 mM phosphate, 1 mM EDTA, pH 6.8, as eluent. The protein fraction off the column was concentrated to a volume of approximately 2 mL in a speed-vac. The protein solution was then mixed with 5'-32P-labeled 508.CF-3'-maleimide (8.9 nmole) and the resulting solution was stirred at room temperature under nitrogen for 16 hours. The reaction was then quenched by the addition of 600 nmole of bromoacetic acid. After stirring for an additional 2 hours at room temperature, the reaction mixture was concentrated to a volume of about 0.5 mL by centrifugal ultrafiltration. A mixture of the conjugate and unconjugated protein was separated from unreacted 508.CF-maleimide by native Polyacrylamide Gel Electrophoresis (PAGE) on a 7.5% density gel (3% cross-linked) in TBE buffer, pH 8, run at constant current of 10 mA at 4 deg C. The mixture of the conjugate and the unconjugated protein was eluted from the gel matrix by a crush and soak method in PBS buffer (138 mM NaCl, 8.1 mM Na₂HPO₄, 2.7 mM KCl, 1.2 mM KH₂PO₄, pH 7.4).

[0108] To further separate the conjugate which incorporates the oligonucleotide, from the unconjugated protein, an affinity purification is called for. A typical example of separating the conjugate with the incorporated oligonucleotide from the contaminating unconjugated protein is the utilization of a solid phase (e.g. functionalized polystyrene or controlled pore glass) immobilized with the complementary oligonucleotide. By performing a hybridization, separation, and washing steps, one can capture the desired conjugate onto the solid phase, wash away the unconjugated protein, and continue on with a releasing step to regain the purified conjugate from the solid phase. Such purification/enrichment steps can be easily done by those skilled in the arts of nucleic acid hybridization.

Example 13.

Preparation of antibody-DNA (anti-TSH-508.CF) conjugate

[0109] This example further illustrates the range of uses for the compounds of the instant invention. In this example antibody-DNA conjugate suitable for carrying a large number of chemiluminescent tags is generated. The conjugate can be prepared in three steps using an alternative thiol-maleimido coupling chemistry.

Preparation of thiolated anti-TSH and conjugation to maleimide-508.CF:

[0110] A solution of anti-TSH (20 mg, 133 nmoles) in 2 mL 0.2 M phosphate buffer, pH 8, containing 150 mM NaCl and 5 mM EDTA was treated with 2-iminothiolane (Traut's reagent, 0.37 mg, 20 equivalents) under nitrogen. The reaction was stirred at room temperature for 1 hour under nitrogen and the thiolated protein was isolated by size-exclusion chromatography on a column of Sephadex G-25 (3.7 x 42 cm) using 10 mM phosphate, pH 6.8, containing 0.5 mM EDTA as eluent. The protein fraction eluting off the column (yield 17 mg, 113 nmoles) was concentrated under reduced pressure in a speed-vac to a volume of approximately 2 mL. This solution was then mixed with maleimide-508.CF (12 nmoles). The reaction was stirred at room temperature for 24 hours under nitrogen by which time the solution had turned slightly cloudy.


[0111] The reaction mixture was then loaded on a column of DEAE-cellulose (1 x 15 cm) equilibrated with 20 mM phosphate, pH 7. The column was eluted with a gradient of 0 to 200 mM NaCl in 20 mM phosphate, pH 7, followed by 200 mM NaCl in 20 mM phosphate, pH 7, to elute excess unreacted protein. Elution with 500 mM NaCl in the same buffer, eluted a mixture of the conjugate and unreacted DNA. Following concentration of the high salt fraction off the ion-exchange column by ultrafiltration, separation of the conjugate from unreacted DNA was achieved by size-exclusion chromatography on a column of Sephadex G-75 (1.5 x 40 cm) using water as eluent at 0.5 mL/min. The solution of the conjugate isolated by this procedure was lyophilized to dryness.

[0112] Characterization of the conjugate and calculation of conjugate yield was performed by UV spectrophotometry. The UV spectrum of the conjugate isolated by the procedure described above was identical in appearance to that of a 1:1 mixture of 508.CF and anti-TSH. This UV spectrum was characterized by an absorption maximum at -270 nm and a 1:1 correspondence between the A_{260} and A_{280} values. The value of A_{260} and A_{280} in the UV spectrum of the conjugate was increased by a factor of 1.5 when compared with DNA alone and protein alone respectively. From this result, the yield of the 1:1 conjugate of 508.CF and anti-TSH was calculated to be 1.6 nmoles (13% overall from maleimide 508.CF).

[0113] The conjugates thus produced can be covalently linked to MHS-DHAE chemiluminescent label by the previously described procedures, among others.

Claims

1. A hydrophilic acridinium ester **characterised in that** it corresponds to the following general formula:









wherein R_1 represents sulfopropyl or sulfoethyl; R_2 represents hydrogen, methoxy, ethoxy, nitro or halogen; R_5 and R_7 represent hydrogen; R_4 and R_3 independently represent methyl, ethyl or isopropyl; and R_6 represents N-succinimidylloxycarbonyl;


and

wherein R_5 , R_6 and R_7 substituent positions on the phenoxy ring are interchangeable.

2. An acridinium ester as claimed in claim 1 wherein R_1 represents sulfopropyl; R_2 , R_5 and R_7 represent hydrogen; R_4 and R_3 represent methyl; and R_6 represents N-succinimidylloxycarbonyl.

3. An acridinium ester conjugate **characterised in that** it comprises an acridinium ester as claimed in claim 1 or claim 2 covalently bonded directly or via a bifunctional linker selected from hexyl-1,6-diamine, ethylene diamine or aminocaproic acid to a small biologically active molecule or a macro-biomolecule.
4. A conjugate as claimed in claim 3 wherein an oligonucleotide is conjugated to the macro-biomolecule, which is further conjugated to multiple acridinium esters and optionally multiple hydrophilic polymers.
5. A conjugate as claimed in claim 4 wherein the oligonucleotide is a gene probe.
6. A conjugate as claimed in claim 3 wherein the macro-biomolecule is selected from protein, peptide, inactivated protein, DNA, RNA, oligonucleotide, neurotransmitter, hormone, steroid hormone, virus, bacterium, toxin and cytokine, or the small biologically active molecule is a hapten.
7. A conjugate as claimed in claim 6 wherein the virus is Rubella virus, or the protein is an antibody, antibody fragment, avidin, streptavidin, allergen, receptor protein, DNA binding protein, neurotransmitter, viral antigen or bacterial antigen or the hapten is a steroid hormone.
8. A conjugate as claimed in claim 3 wherein the small biologically active molecule is a steroid hormone which is testosterone, and the bridge and linker arm are connected by a C-19-C linkage, olefinic C-19 linkage or C19-O linkage, or wherein the small biologically active molecule is a steroid hormone which is testosterone, and the conjugation occurs by hexyl-1,6-diamine, ethylene diamine or aminocaproic acid, or wherein the small biologically active molecule is a steroid hormone which is a testosterone derivative selected from:





wherein A represents a functionalized group selected from -OH, -N(R₁₃)₂, -NHR₁₃, nitro, halide, -OR₁₃, -OCOR₁₃, -tosylate, -mesylate, -azide, -isocyanate, -thioisocyanate, -SR₁₃, -SH, -SSR₁₃, -phosphonate, -sulfonate, -NHCOOR₁₃, -NHCOR₁₃, -NHCONH₂, hydrazine, cyanide, and -R₁₃; wherein R₁₃ represents alkyl, alkenyl, alkynyl or aralkyl having up to 10 carbon atoms and up to 5 heteroatoms; B represents -(CH₂)_n wherein n represents an integer 1-4 or -C=, A is preferably omitted; and X represents carboxylate, -COOR₁₂, -CONHR₁₂, wherein R₁₂ represents alkyl, alkenyl, alkynyl, aryl or aralkyl, having up to 15 carbon atoms and up to 5 heteroatoms, or carbonyl attached with proper leaving groups.

9. A conjugate as claimed in claim 8 wherein the said leaving groups are halide, N-hydroxysuccinimide or imidazole.

10. A method for measuring the amount of analyte in a sample **characterised in that** it comprises:

- (a) detecting an analyte specifically and in proportion to its concentration in a sample; and
- (b) measuring a signal which is generated by an acridinium ester as claimed in claim 1 that is directly or indirectly proportional to the concentration of an analyte in a sample.

11. A method as claimed in claim 10 wherein the detecting comprises:

- (a) contacting a sample with an acridinium ester labelled detector molecule, the said detector molecule optionally being a complex of two or more molecular entities;
- (b) sequestering bound detector and analyte;
- (c) washing away excess detector; and
- (d) measuring a signal from an analyte bound detector;

or comprises:

- (a) contacting a sample with an acridinium ester labelled competitive tracer and a specific binder for an analyte;
- (b) retrieving the specific binder; and
- (c) determining a signal generated by the bound tracer;

or comprises:

- (a) contacting a sample with an acridinium ester labelled competitive tracer and a specific binder for an analyte;
- (b) retrieving the specific binder; and
- (c) determining a signal generated by the unbound tracer;

or comprises:

- (a) contacting a sample with a first specific binder and a second acridinium ester labelled specific binder; and
- (b) detecting a signal;


or comprises:

- (a) contacting a sample with an acridinium ester labelled detector and a competitive binder to the detector;
- (b) sequestering the detector bound to the competitive binder;
- (c) washing away excess detector and the detector bound analyte;
- and
- (d) measuring a signal from the detector bound to the competitive binder;

a releasing agent optionally being used.

Patentansprüche

1. Hydrophiler Acridiniumester,
dadurch gekennzeichnet, daß
er der nachfolgenden allgemeinen Formel entspricht:



wobei R₁ Sulfopropyl oder Sulfoethyl darstellt; R₂ Wasserstoff, Methoxy, Ethoxy, Nitro oder Halogen darstellt; R₃, R₅ und R₇ Wasserstoff darstellen; R₄ und R₈ unabhängig Methyl, Ethyl oder Isopropyl darstellen; und R₆ N-Succinimidylloxycarbonyl darstellt;


und

wobei die R₅-, R₆- und R₇-Substituentenpositionen an dem Phenoxyring austauschbar sind.

2. Acridiniumester nach Anspruch 1, wobei R₁ Sulfopropyl darstellt; R₂, R₃, R₅ und R₇ Wasserstoff darstellen; R₄ und R₈ Methyl darstellen; und R₆ N-Succinimidylloxycarbonyl darstellt.
3. Acridiniumester-Konjugat,
dadurch gekennzeichnet, daß
es einen Acridiniumester wie in Anspruch 1 oder 2 beansprucht umfasst, welcher kovalent direkt oder indirekt über einen bifunktionellen Linker, ausgewählt aus Hexyl-1,6-diamin, Ethylendiamin oder Aminocapronsäure an ein kleines biologisch aktives Molekül oder ein Makro-Biomolekül gebunden ist.
4. Konjugat nach Anspruch 3, wobei ein Oligonukleotid an das Makro-Biomolekül konjugiert ist, welches weiterhin an mehrfache Acridiniumester und wahlweise mehrfache hydrophile Polymeren konjugiert ist.
5. Konjugat nach Anspruch 4, wobei das Oligonukleotid eine Gensonde ist.
6. Konjugat nach Anspruch 3, wobei das Makro-Biomolekül ausgewählt ist aus einem Protein, Peptid, inaktivierten Protein, DNA, RNA, Oligonukleotid, Neurotransmitter, Hormon, Steroidhormon, Virus, Bakterium, Toxin und Cytokin, oder das kleine biologisch aktive Molekül ein Hapten ist.
7. Konjugat nach Anspruch 6, wobei das Virus das Rubella-Virus ist, oder das Protein ein Antikörper ist, ein Antikör-

perfragment, Avidin, Streptavidin, ein Allergen, ein Rezeptorprotein, ein DNA-bindendes Protein, ein Neurotransmitter, ein virales Antigen oder ein bakterielles Antigen, oder das Hapten ein Steroidhormon ist.

8. Konjugat nach Anspruch 3, wobei das kleine biologisch aktive Molekül ein Steroidhormon ist, welches Testosteron ist, und der Brücken- und Linker-Arm durch eine C-19-C-Verbindung, eine olefinische C-19-Verbindung oder C-19-O-Verbindung verbunden sind, oder wobei das kleine biologisch aktive Molekül ein Steroidhormon ist, welches Testosteron ist, und die Konjugation durch Hexyl-1,6-diamin, Ethylendiamin oder Aminocaprinsäure erfolgt, oder wobei das kleine biologisch aktive Molekül ein Steroidhormon ist, welches ein Testosteronderivat ist, ausgewählt aus:





15

- 25

- 30

- 35

- 40

- 45

- 50

55

- 28

- a) In-Kontakt-bringen einer Probe mit einem ersten spezifischen Binder und einem zweiten, Acridiniumester-markierten spezifischen Binder; und
- b) Detektieren eines Signals;


oder, der Nachweis umfasst:

- a) In-Kontakt-bringen einer Probe mit einem Acridiniumester-markierten Detektor und einem zu dem Detektor kompetitiven Binder,
- b) Maskieren des Detektors, der an den kompetitiven Binder gebunden ist;
- c) Abwaschen von überschüssigem Detektor und des Detektor-gebundenen Analyten; und
- d) Messen eines Signals von dem an den kompetitiven Binder gebundenen Detektor,

wobei wahlweise ein Trennmittel verwendet wird.

Revendications

1. Un ester d'acridinium hydrophile **caractérisé en ce qu'il** correspond à la formule générale suivante :










dans laquelle R_1 représente un groupe sulfopropyle ou sulfoéthyle ; R_2 représente l'hydrogène, un groupe méthoxy, éthoxy, nitro ou un halogène ; R_3 , R_5 et R_7 représentent de l'hydrogène ; R_4 et R_8 représentent indépendamment un groupe méthyle, éthyle ou isopropyle ; et R_6 représente un groupe N-succinimidyloxycarbonyl ; et dans laquelle les positions des substituants R_5 , R_6 et R_7 sur le noyau phénoxy sont interchangeables.

2. Un ester d'acridinium tel que revendiqué dans la revendication 1, dans lequel R_1 représente un groupe sulfopropyle ; R_2 , R_3 , R_5 et R_7 représentent de l'hydrogène ; R_4 et R_8 représentent des groupes méthyle ; et R_6 représente un groupe N-succinimidyloxycarbonyl.
3. Un conjugué d'ester d'acridinium **caractérisé en ce qu'il** comprend un ester d'acridinium tel que revendiqué dans la revendication 1 ou la revendication 2, lié par covalence, directement ou indirectement par l'intermédiaire d'un segment de liaison bifonctionnel choisi parmi l'hexyl-1,6-diamine, l'éthylènediamine ou l'acide aminocaproïque, à une petite molécule biologiquement active ou une macromolécule biologique.
4. Un conjugué tel que revendiqué dans la revendication 3, dans lequel un oligonucléotide est conjugué à la macromolécule biologique, qui est en encore conjuguée à des esters d'acridinium multiples et facultativement des polymères hydrophiles multiples.
5. Un conjugué tel que revendiqué dans la revendication 4, dans lequel l'oligonucléotide est une sonde génétique.
6. Un conjugué tel que revendiqué dans la revendication 3, dans lequel la macromolécule biologique est choisie parmi une protéine, un peptide, une protéine inactivée, un ADN, un ARN, un oligonucléotide, un neurotransmetteur,

une hormone, une hormone stéroïde, un virus, une bactérie, une toxine et une cytokine, ou bien la petite molécule biologiquement active est un haptène.

7. Un conjugué tel que revendiqué dans la revendication 6, dans lequel le virus est le virus de la rubéole, ou bien la protéine est un anticorps, un fragment d'anticorps, l'avidine, la streptavidine, un allergène, un récepteur protéique, une protéine liant l'ADN, un neurotransmetteur, un antigène viral ou un antigène bactérien, ou bien l'haptène est une hormone stéroïde.
8. Un conjugué tel que revendiqué dans la revendication 3, dans lequel la petite molécule biologiquement active est une hormone stéroïde qui est la testostérone et le pont ou segment de liaison est attaché par une liaison C à C-19, une liaison oléfinique à C-19 ou une liaison O à C-19, ou dans lequel la petite molécule biologiquement active est une hormone stéroïde qui est la testostérone et la conjugaison a lieu par l'hexyl-1,6-diamine, l'éthylènediamine ou l'acide aminocaproïque, ou dans lequel la petite molécule biologiquement active est une hormone stéroïde qui est un dérivé de testostérone choisi parmi :



où A représente un groupe fonctionnalisé choisi parmi -OH, -N(R₁₃)₂, -NHR₁₃, nitro, halogénure, -OR₁₃, -OCOR₁₃, -tosylate, -mésylate, -azide, -isocyanate, -thioisocyanate, -SR₁₃, -SH, -SSR₁₃, -phosphonate, -sulfonate, -NHCOOR₁₃, -NHCOR₁₃, -NHCONH₂, hydrazine, cyanure et -R₁₃ ; où R₁₃ représente un groupe alkyle, alcényle, alcynyle ou aralkyle ayant jusqu'à 10 atomes de carbone et jusqu'à 5 hétéroatomes ; B représente -(CH₂)_n - où n représente un nombre entier de 1 à 4, ou -C= ; lorsque B représente -C=, A est de préférence omis ; et X représente un groupe carboxylate, -COOR₁₂, -CONHR₁₂, où R₁₂ représente un groupe alkyle, alcényle, alcynyle, aryle ou aralkyle ayant jusqu'à 15 atomes de carbone et jusqu'à 5 hétéroatomes, ou carbonyle attaché par des groupes partants appropriés.

9. Un conjugué tel que revendiqué dans la revendication 8, dans lequel lesdits groupes partants sont un halogénure, le N-hydroxysuccinimide ou l'imidazole.

10. Un procédé pour mesurer la quantité d'un analyte dans un échantillon, **caractérisé en ce qu'il comprend** :

- (a) la détection d'un analyte de façon spécifique et proportionnelle à sa concentration dans un échantillon ; et
- (b) la mesure d'un signal qui est engendré par un ester d'acridinium tel que revendiqué dans la revendication 1, qui est directement ou indirectement proportionnel à la concentration d'un analyte dans un échantillon.

11. Procédé tel que revendiqué dans la revendication 10, dans lequel la détection consiste à :

- (a) mettre en contact un échantillon avec une molécule de détecteur marquée par un ester d'acridinium, ladite molécule de détecteur étant facultativement un complexe de deux ou plusieurs entités moléculaires ;
- (b) séquestrer le détecteur et l'analyte liés ;
- (c) éliminer l'excès de détecteur par lavage ; et
- (d) mesurer un signal émanant du détecteur lié à l'analyte ;

ou consiste à :

- (a) mettre en contact un échantillon avec un traceur compétitif marqué par un ester d'acridinium et un liant spécifique pour un analyte ;
- (b) récupérer le liant spécifique ; et
- (c) déterminer un signal engendré par le traceur lié ;

ou consiste à :

- (a) mettre en contact un échantillon avec un traceur compétitif marqué par un ester d'acridinium et un liant spécifique pour un analyte ;
- (b) récupérer le liant spécifique ; et
- (c) déterminer un signal engendré par le traceur non lié ;

ou consiste à :

- (a) mettre en contact un échantillon avec un premier liant spécifique et un second liant spécifique marqué par un ester d'acridinium ; et
- (b) détecter un signal ;

ou consiste à :

- (a) mettre en contact un échantillon avec un détecteur marqué par un ester d'acridinium et un liant compétitif pour le détecteur ;
- (b) séquestrer le détecteur lié au liant compétitif ;
- (c) éliminer par lavage l'excès de détecteur et l'analyte lié au détecteur ; et
- (d) mesurer un signal émanant du détecteur lié au liant compétitif ;

un agent de libération étant facultativement utilisé.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6

Second Antibody Solid Phase: Competitive



Polyclonal Rb α Testosterone

Figure 7

Competitive Hapten Solid Phase Format





Figure 8

ACS Specific IgE Assay

Allergen-Biotin:Streptavidin-AE detector



IgE Anti-Allergen

Figure 9